



Faculty of Resource Science and Technology

**Phenotypic, Taxonomy, Phytochemical and Physiological Characterization
of Black Pepper (*Piper nigrum* L.) Cultivars in Malaysia**

Chen Yi Shang

**Doctor of Philosophy
2020**

Phenotypic, Taxonomy, Phytochemical and Physiological Characterization of
Black Pepper (*Piper nigrum* L.) Cultivars in Malaysia

Chen Yi Shang

A thesis submitted

In fulfilment of the requirements for the degree of Doctor of Philosophy

(Plant Science)

Faculty of Resource Science and Technology
UNIVERSITI MALAYSIA SARAWAK
2020

DECLARATION

I declare that the work in this thesis was carried out in accordance with the regulation of Universiti Malaysia Sarawak. Except where due acknowledgments have been made, the work is that of the author alone. The thesis has not been accepted for any degree and is not concurrently submitted in candidature of any other degree.



.....

Signature

Name: Chen Yi Shang

Matic No.: 16010129

Faculty of Resource Science and Technology

Universiti Malaysia Sarawak

Date:

ACKNOWLEDGEMENT

My greatest gratitude goes to my supervisor, Prof. Dr. Cheksum Tawan and co-supervisor, Dr. Mohd Razip Bin Asaruddin and Assoc. Prof. Dr. Hairul Azman, for their constant guidance and continuous encouragement throughout the study. Their valuable comments and outlook gave a broader perspective in my research career that will help me forever in my plant research. Heartfelt thanks and appreciations are extended to Dr. Maclin Dayod, section head of Smart Farming Section, Agricultural Research Centre Semongok (ARC), Sarawak for allowing me to use planthouse facilities at ARC. Not forgetting Mr. Kevin Muiyang Tawie Anak Sulok, Research Officer (Plant Physiologist), Malaysian Pepper Board for his assistance in data collection via Photosynthesis System and LAI Analyzer for physiological study. Appreciation also goes to Miss. Nurasyikin Bt. Abdul Rahman, Research Officer (chemist) that assists in interpretation of phytochemical analysis. My deepest appreciation also goes to Mr. Wan Anak Ambi, Mr. Mohamad Khairul Nizam Bin Bakri and Mr. Juvian Jacob for their continuous effort in maintaining the trial plot and assistance in data collection.

This project is fully funded by Malaysian Pepper Board. Fund is sourced from Economic Planning Unit (EPU) Malaysia, under Malaysian Pepper Board RMk-11 Development Project entitled 'Projek Perneriksaan Sumber Manusia'. My acknowledgement is also extended to Human Resource Department, Malaysian Pepper Board for the study fees sponsorship.

DEDICATION

The infinite supports from my family and friends will never be forgotten. I sincerely express my gratitude to my beloved parents, Mr. Chen Mau Kiun and Mdm. Chai Ah Na for their unequivocal sacrifices to ascertain better life and education for me. I am also sincerely thankful to my wife, Mdm. Liew Nyuk Chin and my children (Ashton, Ryan and Davis) for giving moral support along my study. Endless encouragements and love from all of my family members are the strength for me to move on and enable me to accomplish this dissertation successfully.

ABSTRACT

To improve the global market competitiveness of Malaysian black pepper, the quality of peppercorn needs to be strengthened. The varietal regulation at farm level is the prerequisite to enhance quality of peppercorn toward strengthening the marketing strategy. This project has comprehensively documented the morphology of ten important black pepper cultivars in Malaysia, namely ‘Semongok Aman’, ‘Kuching’, ‘Semongok Emas’, ‘Semongok Perak’, ‘Semongok 1’, ‘Nyerigai’, ‘India’, ‘Lampung Daun Lebar’, ‘Sarikei’ and ‘Yong Petai’. The key diagnosis characteristics are ‘Semongok Aman’ (Mucronate shaped leaf apex; High percentage of fruit set high, *ca.* 76%), ‘Kuching’ (High number of inflorescence (spike) per branch per node, *ca.* 58.67; High number of node/feet of stem, *ca.* 4.73), ‘Semongok Emas’ (Leaf colour of Green Group 137 Series; Fruit colour of Green Group 141 Series), ‘Semongok Perak’ (Low conversion rate for production of dried black peppercorn, *ca.* 36.12%; Low conversion rate for production of dried white peppercorn, *ca.* 24.21%), ‘Semongok 1’ (Cordate shaped leaf; Anthocyanin absent at shoot tip), ‘Nyerigai’ (Erect branching behaviour at mature vine stage), ‘India’ (Lanceolate shaped leaf; Light seed weight, *ca.* 0.14 g), ‘Lampung Daun Lebar’ (Oblique shaped leaf base; Blade length-width ratio low, *ca.* 1.52), ‘Sarikei’ (Small leaf area, *ca.* 36.90 cm²; Short inflorescence, *ca.* 6.06 cm; Short fruit spike, *ca.* 8.07 cm; Small fruit size, *ca.* 5.78 mm; Small seed size, *ca.* 3.84 mm; Thin pericarp, *ca.* 1.73 mm) and ‘Yong Petai’ (Long inflorescence length, *ca.* 12.75 cm; Long fruit spike length, *ca.* 17.07 cm; Thin inflorescence, *ca.* 2.90 mm). Phenetic analysis and Principle Component Analysis enlightened that ‘Semongok Aman’ and ‘Semongok 1’ have high distinctive value for identification while ‘Kuching’ and ‘Sarikei’ have high similarity, thus identification are more complicated. In phytochemical study, analysis showed ‘Semongok Aman’ and ‘Lampung Daun Lebar’ has higher piperine

(4.20% and 4.25% respectively) and essential oil content (2.50% and 2.24% respectively). At the same time, this analysis also proved the high pleasant aroma index of peppercorn of ‘Nyerigai’ (26.11), ‘Sarikei’ (25.20) and ‘Kuching’ (18.75). In physiological analysis where the evaluation is based on gas exchange performance and leaf area index, ‘Kuching’ was proven most productive, followed by ‘Nyerigai’ and ‘India’. Besides, the efficiency of artificial pollination has been improved also via floral biology study. This study suggests pollen collection for all the ten cultivars, ideally between 0400 h and 1400 h while the stigma was proven more receptive at stage where elongation and spreading of stigmata has occurred and the emergence and wide spreading of stigmata has been completed. In general, this study has ensured efficient *in situ* identification of black pepper cultivar base on the identification key produced. Besides, the commercial potential of each cultivar particularly in pharmacological and fragrance industry has also been identified. It was hoped the study outcome could be used as industry guideline to ensure sustainable development of Malaysian black pepper industry in future.

Keywords: Black pepper cultivar; floral biology study; Malaysia; morphological analysis; phytochemical study; physiological analysis; *Piper nigrum* L.

Pencirian Fenotip, Taxonomi, Fitokimia dan Fisiologi bagi Kultivar Lada Hitam (Piper Nigrum L.) di Malaysia

ABSTRAK

Untuk meningkat persaingan pasaran sedunia lada di Malaysia, kualiti lada perlu dipertingkatkan. Pengawalan varieti lada pada peringkat kebun merupakan langkah awal dalam menjamin kualiti lada seterusnya memperkasa strategi pemasaran lada hitam Malaysia. Projek ini telah mendokumentasi morfologi bagi sepuluh jenis kultivar lada utama di Malaysia secara menyeluruh, iaitu merangkumi 'Semongok Aman', 'Kuching', 'Semongok Emas', 'Semongok Perak', 'Semongok I', 'Nyerigai', 'India', 'Lampung Daun Lebar', 'Sarikei', dan 'Yong Petai'. Kekunci diagnosis bagi ciri-ciri morfologi adalah 'Semongok Aman' (Berbentuk mukronat pada puncak daun dan asas daun; Peratusan kejadian buah yang tinggi, kira-kira 76%), 'Kuching' (Bilangan tangkai bunga yang tinggi, kira-kira 58.67; Bilangan buku ruas dalam sekaki batang yang tinggi, kira-kira 4.73), 'Semongok Emas' (Warna daun jenis Green Group 137 Series); Warna buah jenis Green Group 141 Series), 'Semongok Perak' (Peratus pertukaran lada hitam yang rendah, kira-kira 36.12%; Peratus pertukaran lada putih yang rendah, kira-kira 24.21%), 'Semongok I' (Daun berbentuk kordat; Antosianin tidak hadir pada hujung pucuk), 'Nyerigai' (Cabang sisi bersifat menegak pada peringkat dewasa pokok), 'India' (Daun berbentuk lanseolat; bijih lada ringan, kira-kira 0.14 g), 'Lampung Daun Lebar' (Asas daun berbentuk serong; Nisbah panjang-lebar daun rendah, kira-kira 1.52), 'Sarikei' (Keluasan daun kecil, kira-kira, 36.90 cm²; Tangkai bunga pendek, kira-kira 6.06 cm; Tangkai buah pendek, kira-kira 8.07 cm; Buah bersaiz kecil, kira-kira 5.78 mm; Bijih lada bersaiz kecil, kira-kira 3.84 mm; Kulit buah nipis, kira-kira 1.73 mm) dan 'Yong Petai' (Tangkai bunga panjang, kira-kira 12.75 cm; Tangkai buah panjang, kira-kira 17.07 cm;

Tangkai bunga nipis, kira-kira 2.90 mm). Analisis finetik dan Analisis Prinsip Komponen telah membuktikan 'Semongok Aman' dan 'Semongok I' mempunyai nilai perbezaan yang tinggi untuk pengecaman manakala 'Kuching' dan 'Sarikei' mempunyai persamaan yang tinggi, jadi pengecaman lebih merumitkan. Dalam kajian fitokimia, analisis menunjukkan 'Semongok Aman' dan 'Lampung Daun Lebar' mempunyai kandungan piperine (4.20% dan 4.25% masing-masing) dan kandungan minyak pati (2.50% dan 2.24% masing-masing) yang tinggi. Pada masa yang sama, analisis ini juga membuktikan index aroma wangi yang tinggi pada bijih lada yang dihasilkan oleh 'Nyerigai' (26.11), 'Sarikei' (25.20) dan 'Kuching' (18.75). Dalam analisis fisiologi yang dinilai berdasarkan prestasi pertukaran gas dan index keluasan daun, 'Kuching' telah dibuktikan paling produktif, diikuti oleh 'Nyerigai' dan 'India'. Selain itu, keberkesanan pendebungaan artifisial juga dipertingkatkan melalui kajian biologi bunga. Kajian ini mencadangkan pengumpulan polen bagi semua kultivar sebaiknya dijalankan di antara 0400 jam dan 1400 jam manakala stigma dibuktikan lebih berdaya terima pada peringkat pemanjangan dan perkembangan stigmata telah bermula dan pada peringkat kemunculan dan perkembangan stigmata telah lengkap. Secara umumnya, kajian ini telah memastikan keberkesanan pengecaman kultivar lada hitam secara in situ. Selain itu, potensi komersial bagi setiap kultivar juga dikenalpasti, khususnya dalam industri farmakologikal dan pewangi. Diharapkan hasil kajian ini boleh dijadikan sebagai panduan industri, untuk memastikan kemapanan pembangunan industri lada hitam di Malaysia pada masa depan.

Kata kunci: Kultivar lada hitam; kajian biologi bunga; analisis morfologi; kajian fitokimia; analisis fisiologi; *Piper nigrum L.*; Malaysia.

TABLE OF CONTENTS

	Page
DECLARATION	i
ACKNOWLEDGEMENT	ii
DEDICATION	iii
ABSTRACT	iv
<i>ABSTRAK</i>	vi
TABLE OF CONTENTS	viii
LIST OF TABLES	xvi
LIST OF FIGURES	xvi
CHAPTER 1: INTRODUCTION	1
1.1 Background	1
1.2 Problem Statements	4
1.3 Objectives	5
1.4 Organization of the Thesis	5
CHAPTER 2: LITERATURE REVIEW	8
2.1 Background of Black Pepper (<i>Piper nigrum</i> L.)	8
2.1.1 Taxonomic History	8
2.1.2 Taxonomic Description	9
2.2 Commercial Black Pepper Farm in Malaysia	11
2.3 Black Pepper Cultivars	15
2.3.1 Diversity	15

2.3.2 Black Pepper Germplasm in Malaysia	17
2.3.2.1 Accessions Record	17
2.3.2.2 Genetic Relatedness of Black Pepper in Malaysia	18
2.3.3 Quantitative and Qualitative Characteristics to Differentiate Morphology of Black Pepper Cultivars	20
2.4 Phytochemical of Black Pepper	22
2.4.1 Gas Exchange Parameters	22
2.4.2 Importance: Pungency, Aromatic and Medicinal Values	25
2.5 Physiology of Black Pepper Vine	27
2.5.1 Gas Exchange Parameters in Relation with Plant Productivity	27
2.5.2 Leaf Area Index (LAI) of Black pepper	29
2.6 Floral Biology of Black Pepper	31
2.6.1 Flower Morphology and Development	31
2.6.2 Inflorescence Morphology and development	32
2.6.3 Fruit Set and Development	33
2.6.4 Pollination Biology of Black Pepper	36
2.6.4.1 Anther Dehiscence, Stigma Receptivity and Pollen Viability	36
2.6.4.2 Pollination Ecology	37
CHAPTER 3: TAXONOMY OF <i>PIPER NIGRUM</i> L. AND ITS CULTIVARS IN MALAYSIA	40
3.1 Introduction	40
3.2 Materials and Methods	42
3.3 General Botanical Descriptions of <i>Piper nigrum</i> L.	47
3.4 Cultivars of Black Pepper In Malaysia	51

3.4.1 Diversity	51
3.4.1.1 Cultivar ‘Semongok Aman’	52
3.4.1.2 Cultivar ‘Kuching’	55
3.4.1.3 Cultivar ‘Semongok Emas’	58
3.4.1.4 Cultivar ‘Semongok Perak’	61
3.4.1.5 Cultivar ‘Semongok 1’	64
3.4.1.6 Cultivar ‘Nyerigai’	67
3.4.1.7 Cultivar ‘India’	70
3.4.1.8 Cultivar ‘Lampung Daun Lebar’	73
3.4.1.9 Cultivar ‘Sarikei’	76
3.4.1.10 Cultivar ‘Yong Petai’	79
3.5 Conclusions	82
CHAPTER 4: MORPHOLOGICAL CHARACTERISTICS OF	84
BLACK PEPPER CULTIVARS	
4.1 Introduction	84
4.2 Materials and Methods	86
4.2.1 Analysis of Qualitative and Quantitative Trait Variability	86
4.2.1.1 Material Preparation	86
4.2.1.2 Experiment Design	87
4.2.1.3 Data Analysis	89
4.2.2 Phenetic Analysis	91
4.2.2.1 Material Preparation and Experiment Design	91
4.2.2.2 Data Analysis	92
4.3 Results and Discussion	93

4.3.1 Analysis of Qualitative and Quantitative Trait Variability	93
4.3.1.1 Variability Analysis	93
4.3.1.2 Dichotomous Key for Cultivar Diagnosis	114
4.3.1.3 Yield Potential Study	116
4.3.2 Phenetic Analysis	121
4.3.2.1 Cluster Analysis	121
4.3.2.2 Principle Component Analysis	127
4.4 Conclusions	133
CHAPTER 5: PHYTOCHEMICAL CHARACTERIZATION OF	135
BLACK PEPPER CULTIVARS	
5.1 Introduction	135
5.2 Materials and Methods	137
5.2.1 Materials	137
5.2.2 Sample Preparation and GC Measurement	138
5.2.3 HPLC Analysis of Piperine	139
5.2.4 Statistical Analysis	140
5.3 Results and Discussion	140
5.3.1 Essential Oil Analysis	140
5.3.2 Piperine Analysis	155
5.4 Conclusions	157
CHAPTER 6: PHYSIOLOGICAL AND FLORAL BIOLOGY OF	159
BLACK PEPPER CULTIVARS	
6.1 Introduction	159
6.2 Materials and Methods	161

6.2.1 Physiological Study	161
6.2.1.1 Gas Exchange Evaluation	161
6.2.1.1.1 Material preparation	161
6.2.1.1.2 Experiment Design for Gas Exchange Assessment	162
6.2.1.1.3 Data Analysis	164
6.2.1.2 Leaf Area Index	164
6.2.1.2.1 Materials Preparation	164
6.2.1.2.2 Experiment Design for Leaf Area Index Assessment	164
6.2.1.2.3 Data Analysis	165
6.2.2 Floral Biology Study	165
6.2.2.1 Material Preparation	165
6.2.2.2 Experiment Design and Data analysis	166
6.2.2.2.1 Pollen Morphology Study	166
6.2.2.2.2 Anther Dehiscence	167
6.2.2.2.3 Pollen Viability Study	167
6.2.2.2.4 Stigma Receptivity Study	169
6.3 Results and Discussion	170
6.3.1 Physiological Study	170
6.3.1.1 Gas Exchange Evaluation	170
6.3.1.2 Leaf Area Index (LAI) Analysis	179
6.3.2 Floral Biology Study	182
6.3.2.1 Pollen Morphology Study	182
6.3.2.2 Anther Dehiscence	184

6.3.2.3 Pollen Viability Study	185
6.3.2.4 Stigma Receptivity Study	190
6.4 Conclusion	194
CHAPTER 7: GENERAL DISCUSSIONS, CONCLUSIONS AND RECOMMENDATIONS	196
7.1 General Discussions	196
7.2 Conclusions	201
7.2.1 Taxonomy of <i>Piper nigrum</i> L. and Its Cultivars in Malaysia	201
7.2.2 Morphological Characteristics of Black Pepper Cultivars	201
7.2.3 Physiological Characterization of Black Pepper Cultivars	203
7.2.4 Phytochemical and Floral Biology of Black Pepper Cultivars	203
7.3 Recommendations or Future Prospects	204
REFERENCES	207
APPENDICES	239

LIST OF TABLES

		Page
Table 2.1	Black pepper accessions selected for DAMD analysis.	18
Table 2.2	Piperine therapeutic properties for modern medical applications.	27
Table 2.3	Stages of black pepper flower development.	32
Table 2.4	Stages of black pepper inflorescence development.	33
Table 2.5	Stages of black pepper fruit development.	34
Table 2.6	Fieldwork area coverage and observations.	44
Table 4.1	Morphological characteristic used for diagnosis of cultivar distinctness.	90
Table 4.2	Qualitative and quantitative traits used to diagnose the differences among black pepper cultivars.	94
Table 4.3	Yield potential evaluation for the ten cultivars based on morphological yield attributing characters.	119
Table 4.4	Characteristics and character states used for phenetic analysis of the ten black pepper cultivars.	122
Table 4.5	Data matrix corresponding to Table 4.4.	124
Table 4.6	Total variance obtained using Principle Component Analysis (PCA).	128
Table 4.7	Matrix component value for all the distinguishing characteristics.	130
Table 5.1	The molecular structure and content of major monoterpene,	136

sesquiterpenes and oxygen-containing sesquiterpenes in pepper berries.

Table 5.2	The gradient HPLC for the analysis.	139
Table 5.3	Percent content of essential oil in peppercorn of the ten cultivars.	141
Table 5.4	Essential oil constituent composition of the ten cultivars.	143
Table 5.5	Characteristics and character states of essential oil constituent of the ten cultivars for the PCA analysis.	146
Table 5.6	Data matrix corresponding to Table 5.5.	147
Table 5.7	Total variance obtained using Principle Component Analysis (PCA) for the essential oil constituent analysis.	149
Table 5.8	Matrix component value for all the distinguishing characteristics for the essential oil constituent analysis.	152
Table 5.9	Aroma analysis based on monoterpene constituents in essential oil of the ten cultivars.	154
Table 5.10	Percent content of piperine in peppercorn of the ten cultivars.	156
Table 6.1	LAI for ten black pepper cultivars.	181
Table 6.2	Receptivity difference at three stages of stigma for ten cultivars.	193

LIST OF FIGURES

	Page
Figure 2.1 Big scale black pepper farm located at Sri Aman, Sarawak, Malaysia.	12
Figure 2.2 Vine uniformity contributed by systematic black pepper planting system in Malaysia.	13
Figure 2.3 UPGMA dendrogram of DAMD similarities among 43 black pepper accessions. Sourced from Ho et al. (2005).	19
Figure 3.1 Completely Randomized Design (CRD) pot arrangement for the ten treatments.	43
Figure 3.2 Botanical drawing of cultivated black pepper.	49
Figure 3.3 Photography documentation of important morphological characteristics of cultivated black pepper.	50
Figure 3.4 Photography documentation of cv. ‘Semongok Aman’.	53
Figure 3.5 Dried specimen of cv. ‘Semongok Aman’.	53
Figure 3.6 Distribution of cv. ‘Semongok Aman’.	54
Figure 3.7 Photography documentation of cv. ‘Kuching’.	56
Figure 3.8 Dried specimen of cv. ‘Kuching’.	56
Figure 3.9 Distribution of cv. ‘Kuching’.	57
Figure 3.10 Photography documentation of cv. ‘Semongok Emas’.	59
Figure 3.11 Dried specimen of cv. ‘Semongok Emas’.	59
Figure 3.12 Distribution of cv. ‘Semongok Emas’.	60
Figure 3.13 Photography documentation of cv. ‘Semongok Perak’.	62

Figure 3.14	Dried specimen of cv. ‘Semongok Perak’.	62
Figure 3.15	Distribution of cv. ‘Semongok Perak’.	63
Figure 3.16	Photography documentation of cv. ‘Semongok 1’.	65
Figure 3.17	Dried specimen of cv. ‘Semongok 1’.	65
Figure 3.18	Distribution of cv. ‘Semongok 1’.	66
Figure 3.19	Photography documentation of cv. ‘Nyerigai’.	68
Figure 3.20	Dried specimen of cv. ‘Nyerigai’.	68
Figure 3.21	Distribution of cv. ‘Nyerigai’.	69
Figure 3.22	Photography documentation of cv. ‘India’.	71
Figure 3.23	Dried specimen of cv. ‘India’.	71
Figure 3.24	Distribution of cv. ‘India’.	72
Figure 3.25	Photography documentation of cv. ‘Lampung Daun Lebar’.	74
Figure 3.26	Dried specimen of cv. ‘Lampung Daun Lebar’.	74
Figure 3.27	Distribution of cv. ‘Lampung Daun Lebar’.	75
Figure 3.28	Photography documentation of cv. ‘Sarikei’.	77
Figure 3.29	Dried specimen of cv. ‘Sarikei’.	77
Figure 3.30	Distribution of cv. ‘Sarikei’.	78
Figure 3.31	Photography documentation of cv. ‘Yong Petai’.	80
Figure 3.32	Dried specimen of cv. ‘Yong Petai’.	80
Figure 3.33	Distribution of cv. ‘Yong Petai’.	81
Figure 4.1	Randomized Complete Block Design (RCBD) for the field grown vine of ten cultivars.	88
Figure 4.2	Completely Randomized Design (CRD) pot arrangement for experimental plot under controlled environment.	89

Figure 4.3	Cultivar designation and leaf shape.	99
Figure 4.4	Inflorescence.	100
Figure 4.5	Fruit spike.	100
Figure 4.6	Cross section of ripe and mature fruit.	101
Figure 4.7	Shoot tips.	101
Figure 4.8	Leaf length-width ratio.	102
Figure 4.9	Number of flower per inflorescence.	103
Figure 4.10	Leaf width (cm).	103
Figure 4.11	Leaf area (cm ²).	104
Figure 4.12	Spike or inflorescence thicknesses (mm).	104
Figure 4.13	Number of inflorescence (spike) per branch per node.	105
Figure 4.14	Percent fruit set (%).	105
Figure 4.15	Fruit weight (single berry) (g).	106
Figure 4.16	Internode lengths (cm).	106
Figure 4.17	Number of nodes per foot of stem.	107
Figure 4.18	Leaf length (cm).	107
Figure 4.19	Inflorescence lengths (cm).	108
Figure 4.20	Fresh to dried (white pepper conversion rate %).	109
Figure 4.21	Pericarp thicknesses (mm).	109
Figure 4.22	Seed diameter (mm).	110
Figure 4.23	Single fresh berry sizes in diameter (mm).	110
Figure 4.24	Fresh to dried (black pepper conversion rate %).	111
Figure 4.25	Fruit spike length (cm).	111
Figure 4.26	Seed weight (10 ⁻²) (g).	112

Figure 4.27	Phenogram presented based on average linkage (between groups) using Squared Euclidean Distance Method.	125
Figure 4.28	Scree plot showing cut-off point of extracted components for PCA of morphological characteristics.	127
Figure 5.1	Scree plot showing cut-off point of extracted components for PCA of phytochemical analysis.	148
Figure 6.1	Plagiotropic branch of black pepper.	162
Figure 6.2	Photosynthetic performance for ten black pepper cultivars.	171
Figure 6.3	Leaf stomatal conductance evaluation for ten black pepper cultivars.	173
Figure 6.4	Transpiration rate evaluation for ten black pepper cultivars.	175
Figure 6.5	Relationship between net photosynthesis rate and leaf stomatal conductance for the ten black pepper cultivars.	176
Figure 6.6	Relationship between transpiration rate and leaf stomatal conductance for the ten black pepper cultivars.	177
Figure 6.7	Relationship between transpiration rate and net photosynthesis rate for the ten black pepper cultivars.	177
Figure 6.8	Relationship between LAI and net photosynthesis rate in black pepper cultivars.	181
Figure 6.9	SEM micrographs showing size, shape and sculpture of pepper cultivar 'SA' pollen grains.	183
Figure 6.10	Time of anther dehiscence in cultivar 'Semongok Aman' between March and July 2016.	185

Figure 6.11	<i>In vitro</i> germinated pollen of cv. ‘Semongok Aman’ at different viability level.	186
Figure 6.12	Percentage of pollen germination at various stages of pollen for the ten cultivars.	187
Figure 6.13	Pollen tube length at five various stages of the anther dehiscence for the ten cultivars.	189
Figure 6.14	Stages of stigma.	191

CHAPTER 1

INTRODUCTION

1.1 Background

Piper nigrum L., commonly called black pepper is the world renowned spice from the family of *Piperaceaea*. The peppercorn harvested from the plant not only widely used for food preparation that attributed by unique aroma and pungent properties, but also its pharmacological value (Krishnapura, 2009; Menghani et al., 2010; Meghwal & Goswami, 2012; Butt et al., 2013). The alkaloid compound, piperine, is the compound that contributes to the pungency of black pepper (Krishnapura, 2009; Gorgani et al., 2017) while the aroma is attributed to the volatile oil found in the peppercorn (Nirmala Menon & Padmakumari, 2005). The pharmacological potential that has been scientifically proven include immune modulatory, anti-carcinogenic, stimulatory, hepatoprotective and anti-inflammatory by Darshan and Doreswamy (2004); antimicrobial by Yang et al. (2002); antiulcer activities by Bai and Xu (2000); biotransformative effects and enhance the bioavailability of different drugs by Atal et al. (1985) and Wu (2007); protective effect against radiation by Raman and Gaikar (2002b); Increases pancreatic lipase activity and stimulates pancreatic amylase, trypsin, and chymotrypsin by Platel and Srinivasan (2000) and reduction of blood cholesterol, triglycerides, and glucose by Mueller and Hingst (2013).

The cultivar or cultivated variety of black pepper was believed to have derived from the wild one through domestication (Ravindran, 2000) in the center of origin, the

Malabar Island of India. The plant was introduced to Malaysia as early as 1840 by Hakka Chinese (Mullen, 1960) through Babylonian-Chinese sea trade that linked Malabar Coast and South East Asian ports (Ravindran, 2000). Sarawak and Johore region were the main cultivation area at the beginning stage. However, cultivation only focused in Sarawak in later years with more than 98% of country total production was from this state (Eng & Gumbek, 2007). The distribution in peninsula state and Sabah is limited mainly due to the farmer preference to others crop such as oil palm and rubber. Recent total black pepper planting hectare is around 13,000 hectare (Ibragimov et al., 2015), mainly distributed in rural area of Sarawak. Approximately 18,000 to 20,000 pepper farm families, with 68,000 to 70,000 people involved in pepper cultivation (Wong, 2016).

The most common black pepper cultivar is cv. 'Kuching', which is widely planted throughout Sarawak (Sim, 1991). However, Paulus (2007) reported three important cultivars in Malaysia, namely cv. 'Semongok Perak', cv. 'Semongok Aman', and cv. 'Kuching'. Through the International Pepper Community (IPC) exchange program, cv. 'Lampung Daun Lebar' and cv. 'Lampung Daun Kecil' were introduced to Malaysian farmers (Sim, 2007). In recent year, a manual entitled 'Pepper production technology in Malaysia' released by the Malaysian Pepper Board, reported the existence of seven cultivated varieties as common cultivars in Malaysia, including cv. 'Semongok Aman', cv. 'Semongok Emas', cv. 'Kuching', cv. 'Semongok Perak', cv. 'Uthirancotta', cv. 'Nyerigai', and cv. 'PN129'(Paulus, 2011a). However, there is no varietal restriction or regulation in black pepper planting in the past until nowadays. Thus, the diversity of the black pepper cultivar in Malaysia black pepper farm is unrevealed.